

## Computer Graphics

CSE 167 [Win 22], Lecture 10: Curves 2

Ravi Ramamoorthi

<http://viscomp.ucsd.edu/classes/cse167/wi22>

## Outline of Unit


- Bezier curves (last time)
- deCasteljau algorithm, explicit, matrix (last time)
- *Polar form labeling (blossoms)*
- B-spline curves

▪ Not well covered in textbooks (especially as taught here). Main reference will be lecture notes. If you do want a printed ref, handouts from CAGD, Seidel

## Survey Feedback


## Idea of Blossoms/Polar Forms

- (Optional) Labeling trick for control points and intermediate deCasteljau points that makes thing intuitive
- E.g. quadratic Bezier curve  $F(u)$ 
  - Define auxiliary function  $f(u_1, u_2)$  [number of args = degree]
  - Points on curve simply have  $u_1 = u_2$  so that  $F(u) = f(u, u)$
  - And we can label control points and deCasteljau points not on curve with appropriate values of  $(u_1, u_2)$






## Idea of Blossoms/Polar Forms

- Points on curve simply have  $u_1 = u_2$  so that  $F(u) = f(u, u)$
- $f$  is symmetric  $f(0, 1) = f(1, 0)$
- Only interpolate linearly between points with one arg different
  - $f(0, u) = (1-u) f(0, 0) + u f(0, 1)$  Here, interpolate  $f(0, 0)$  and  $f(0, 1) = f(1, 0)$



## Geometric interpretation: Quadratic





### Why Polar Forms?

- Simple mnemonic: which points to interpolate and how in deCasteljau algorithm
- Easy to see how to subdivide Bezier curve (next) which is useful for drawing recursively*
- Generalizes to arbitrary spline curves (just label control points correctly instead of 00 01 11 for Bezier)
- Easy for many analyses (beyond scope of course)



### Subdividing Bezier Curves







Drawing: Subdivide into halves ( $u = \frac{1}{2}$ ) Demo: hw3

- Recursively draw each piece
- At some tolerance, draw control polygon
- Trivial for Bezier curves (from deCasteljau algorithm): hence widely used for drawing

Why specific labels/ control points on left/right?

- How do they follow from deCasteljau?






## Polar Forms: Cubic Bspline Curve

- Labeling little different from in Bezier curve
- No interpolation of end-points like in Bezier
- Advantage of polar forms: easy to generalize




## deCasteljau: Cubic B-Splines

- Easy to generalize using polar-form labels
- Impossible remember without




## deCasteljau: Cubic B-Splines

- Easy to generalize using polar-form labels
- Impossible remember without




## deCasteljau: Cubic B-Splines

- Easy to generalize using polar-form labels
- Impossible remember without



## Explicit Formula (derive as exercise)

$$F(u) = [u^3 \ u^2 \ u \ 1] M \begin{bmatrix} P_0 \\ P_1 \\ P_2 \\ P_3 \end{bmatrix} \quad M = \frac{1}{6} \begin{bmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 0 & 3 & 0 \\ 1 & 4 & 1 & 0 \end{bmatrix}$$



## Summary of HW 3

- BSpline Demo hw3
- Arbitrary number of control points / segments
  - Do nothing till 4 control points (see demo)
  - Number of segments = # cpts - 3
- Segment A will have control pts A, A+1, A+2, A+3
- Evaluate Bspline for each segment using 4 control points (at some number of locations, connect lines)
- Use either deCasteljau algorithm (like Bezier) or explicit form [matrix formula on previous slide]
- Questions?