


Computer Graphics II: Rendering

CSE 168 [Spr 20], Lecture 5: Monte Carlo Integration
Ravi Ramamoorthi

<http://viscomp.ucsd.edu/classes/cse168/sp20>



To Do

- Homework 2 (Direct Lighting) due Apr 24
- Assignment is on edX edge
- START EARLY (NOW)

Motivation


Rendering = integration

- Reflectance equation: Integrate over incident illumination
- Rendering equation: Integral equation

Many sophisticated shading effects involve integrals

- Antialiasing
- Soft shadows
- Indirect illumination
- Caustics

Example: Soft Shadows

$$E(x) = \int_{H^2} L_i(x, \omega) \cos \theta d\omega$$


Challenges

- **Visibility and blockers**
- **Varying light distribution**
- **Complex source geometry**

Source: Agrawala, Ramamoorthi, Heirich, Moll, 2000

Monte Carlo

- Algorithms based on statistical sampling and random numbers
- Coined in the beginning of 1940s. Originally used for neutron transport, nuclear simulations
 - Von Neumann, Ulam, Metropolis, ...
- Canonical example: 1D integral done numerically
 - Choose a set of random points to evaluate function, and then average (expectation or statistical average)

Monte Carlo Algorithms

Advantages

- Robust for complex integrals in computer graphics (irregular domains, shadow discontinuities and so on)
- Efficient for high dimensional integrals (common in graphics: time, light source directions, and so on)
- Quite simple to implement
- Work for general scenes, surfaces
- Easy to reason about (but care taken re statistical bias)


Disadvantages

- Noisy
- Slow (many samples needed for convergence)
- Not used if alternative analytic approaches exist (but those are rare)

Outline

- Motivation
- Overview, 1D integration
- Basic probability and sampling
- Monte Carlo estimation of integrals


Integration in 1D



Slide courtesy of
Peter Shirley

We can approximate

$$\int_0^1 f(x) dx \approx \int_0^1 g(x) dx$$



Standard integration methods like trapezoidal rule and Simpsons rule

Advantages:

- Converges fast for **smooth** integrands
- Deterministic


Disadvantages:

- Exponential complexity in many dimensions
- Not rapid convergence for discontinuities

Slide courtesy of
Peter Shirley

Or we can average


$$\int_0^1 f(x) dx = E(f(x))$$



Slide courtesy of
Peter Shirley

Estimating the average

$$\int_0^1 f(x) dx = \frac{1}{N} \sum_{i=1}^N f(x_i)$$



Monte Carlo methods (randomly choose samples)


Advantages:

- Robust for discontinuities
- Converges reasonably for large dimensions
- Can handle complex geometry, integrals
- Relatively simple implement, reason about

Slide courtesy of
Peter Shirley

Other Domains

$$\int_a^b f(x) dx = \frac{b-a}{N} \sum_{i=1}^N f(x_i)$$




Slide courtesy of
Peter Shirley

Multidimensional Domains

Same ideas apply for integration over ...

- Pixel areas
- Surfaces
- Projected areas
- Directions
- Camera apertures
- Time
- Paths



Outline

- Motivation
- Overview, 1D integration
- *Basic probability and sampling*
- Monte Carlo estimation of integrals

Random Variables

- Describes possible outcomes of an experiment
- In discrete case, e.g. value of a dice roll [$x = 1-6$]
- Probability p associated with each x (1/6 for dice)
- Continuous case is obvious extension

Expected Value

- Expectation Discrete: $E(x) = \sum_{i=1}^n p_i x_i$
Continuous: $E(x) = \int_0^1 p(x)f(x) dx$
- For Dice example:

$$E(x) = \sum_{i=1}^6 \frac{1}{6} x_i = \frac{1}{6} (1+2+3+4+5+6) = 3.5$$

Continuous Probability Distributions

PDF $p(x)$

$$p(x) \geq 0$$

Uniform



CDF $P(x)$

$$P(x) = \int_0^x p(x)dx$$

$$P(x) = \Pr(X < x)$$

$$\Pr(\alpha \leq X \leq \beta) = \int_{\alpha}^{\beta} p(x)dx$$


$$= P(\beta) - P(\alpha)$$

Pat Hanrahan, Spring 2004

Sampling Techniques

Problem: how do we generate random points/directions during path tracing?

- Non-rectilinear domains
- Importance (BRDF)
- Stratified



Generating Random Points

Uniform distribution:

- Use random number generator

Generating Random Points

Specific probability distribution:

- Function inversion
- Rejection
- Metropolis

Common Operations

Want to **sample** probability distributions

- Draw samples distributed according to probability
- Useful for integration, picking important regions, etc.

Common distributions

- Disk or circle
- Uniform
- Upper hemisphere for visibility
- Area luminaire
- Complex lighting like an environment map
- Complex reflectance like a BRDF

Sampling Continuous Distributions

Cumulative probability distribution function

$$P(x) = \Pr(X < x)$$

Construction of samples

Solve for $X = P^{-1}(U)$

Must know:

- The integral of $p(x)$**
- The inverse function $P^{-1}(x)$**

CS348B Lecture 6

Pat Hanrahan, Spring 2004

Generating Random Points

Example: Power Function

Assume

$$p(x) = (n+1)x^n$$

$$P(x) = x^{n+1}$$

$$X \sim p(x) \Rightarrow X = P^{-1}(U) = \sqrt[n+1]{U}$$

Trick

$$Y = \max(U_1, U_2, \dots, U_n, U_{n+1})$$

$$\Pr(Y < x) = \prod_{i=1}^{n+1} \Pr(U_i < x) = x^{n+1}$$

CS348B Lecture 6

Pat Hanrahan, Spring 2004

Sampling a Circle

$$A = \int_0^{2\pi} \int_0^1 r dr d\theta = \int_0^1 r dr \int_0^{2\pi} d\theta = \left(\frac{r^2}{2} \right) \Big|_0^{2\pi} = \pi$$

$$p(r, \theta) dr d\theta = \frac{1}{\pi} r dr d\theta \Rightarrow p(r, \theta) = \frac{r}{\pi}$$

$$p(r, \theta) = p(r)p(\theta)$$

$$p(\theta) = \frac{1}{2\pi}$$

$$P(\theta) = \frac{1}{2\pi}$$

$$\theta = 2\pi U_1$$

$$r = \sqrt{U_2}$$

Pat Hanrahan, Spring 2004

Sampling a Circle

WRONG \neq Equi-Areal

$\theta = 2\pi U_1$
 $r = U_2$

RIGHT = Equi-Areal

$\theta = 2\pi U_1$
 $r = \sqrt{U_2}$

Pat Hanrahan, Spring 2004

Rejection Sampling

Probability

W

Pat Hanrahan, Spring 2004

Rejection Methods

$$I = \int_0^1 f(x) dx$$

$$= \iint_{y < f(x)} dx dy$$

Algorithm

Pick U_1 **and** U_2

Accept U_1 **if** $U_2 < f(U_1)$

Wasteful? Efficiency = Area / Area of rectangle

Pat Hanrahan, Spring 2004

Sampling a Circle: Rejection

```
do {
  X=1-2*U1
  Y=1-2*U2
  while( X^2+ Y^2 >1 )
```

May be used to pick random 2D directions

Circle techniques may also be applied to the sphere


CS348B Lecture 6

Pat Hanrahan, Spring 2004

Outline

- Motivation
- Overview, 1D integration
- Basic probability and sampling
- Monte Carlo estimation of integrals

Monte Carlo Path Tracing



Big diffuse light source, 20 minutes

Motivation for rendering in graphics: Covered in detail in next lecture

Jensen

Monte Carlo Path Tracing




1000 paths/pixel

Jensen

Estimating the average

$\int_0^1 f(x) dx = \frac{1}{N} \sum_{i=1}^N f(x_i)$



Monte Carlo methods (randomly choose samples)

Advantages:

- Robust for discontinuities
- Converges reasonably for large dimensions
- Can handle complex geometry, integrals
- Relatively simple implement, reason about

Slide courtesy of Peter Shirley

Monte Carlo Integration

Definite integral $I(f) \equiv \int_0^1 f(x) dx$

Expectation of f $E[f] \equiv \int_0^1 f(x) p(x) dx$

Random variables $X_i \sim p(x)$
 $Y_i = f(X_i)$

Estimator $F_N = \frac{1}{N} \sum_{i=1}^N Y_i$

Unbiased Estimator

$E[F_N] = I(f)$

$$\begin{aligned} E[F_N] &= E\left[\frac{1}{N} \sum_{i=1}^N Y_i\right] \\ &= \frac{1}{N} \sum_{i=1}^N E[Y_i] = \frac{1}{N} \sum_{i=1}^N E[f(X_i)] \\ &= \frac{1}{N} \sum_{i=1}^N \int_0^1 f(x) p(x) dx \\ &= \frac{1}{N} \sum_{i=1}^N \int_0^1 f(x) dx \\ &= \int_0^1 f(x) dx \end{aligned}$$

Properties

$$\begin{aligned} E\left[\sum_i Y_i\right] &= \sum_i E[Y_i] \\ E[aY] &= aE[Y] \end{aligned}$$


Assume uniform probability distribution for now

CS348B Lecture 6

Pat Hanrahan, Spring 2004

Direct Lighting – Directional Sampling

$E(x) = \int_{\Omega} L(x, \omega) \cos \theta d\omega$



Ray intersection $x^*(x, \omega)$

Sample ω uniformly by Ω

$$Y_i = L(x^*(x, \omega_i), -\omega_i) \cos \theta 2\pi$$

CS348B Lecture 6

Pat Hanrahan, Spring 2004

Direct Lighting - Area Sampling


$E(x) = \int_{\Omega} L_i(x, \omega) \cos \theta d\omega = \int_{A'} L_o(x', \omega') V(x, x') \frac{\cos \theta \cos \theta'}{|x - x'|^2} dA'$

Ray direction $\omega' = x - x'$
Sample x' uniformly by A'

$$Y_i = L_o(x'_i, \omega'_i) V(x, x'_i) \frac{\cos \theta \cos \theta'_i}{|x - x'_i|^2} A$$

$$V(x, x') = \begin{cases} 0 & \text{non-visible} \\ 1 & \text{visible} \end{cases}$$

CS348B Lecture 6 Pat Hanrahan, Spring 2004



Variance

Definition

$V[Y] \equiv E[(Y - E[Y])^2]$
 $= E[Y^2 - 2YE[Y] + E[Y]^2]$
 $= E[Y^2] - E[Y]^2$

Properties

$V[\sum_i Y_i] = \sum_i V[Y_i]$
 $V[aY] = a^2 V[Y]$



Variance decreases with sample size

$V[\frac{1}{N} \sum_{i=1}^N Y_i] = \frac{1}{N^2} \sum_{i=1}^N V[Y_i] = \frac{1}{N} V[Y]$

CS348B Lecture 6 Pat Hanrahan, Spring 2004

Variance for Dice Example?

- Work out on board (variance for single dice roll)



Variance Reduction

Efficiency measure

$$\text{Efficiency} \propto \frac{1}{\text{Variance} \cdot \text{Cost}}$$

Techniques

- Importance sampling
- Sampling patterns: stratified, ...

CS348B Lecture 6 Pat Hanrahan, Spring 2004

Variance Reduction Techniques

- Importance sampling
- Stratified sampling

$$\int_0^1 f(x) dx = \frac{1}{N} \sum_{i=1}^N f(x_i)$$

Importance Sampling

Put more samples where $f(x)$ is bigger

$$\int_{\Omega} f(x) dx = \frac{1}{N} \sum_{i=1}^N Y_i$$

$$Y_i = \frac{f(x_i)}{p(x_i)}$$

Importance Sampling

- This is still unbiased

$$\begin{aligned} E[Y_i] &= \int_{\Omega} Y_i p(x) dx \\ &= \int_{\Omega} \frac{f(x)}{p(x)} p(x) dx \\ &= \int_{\Omega} f(x) dx \end{aligned}$$

for all N

Importance Sampling

- Zero variance if $p(x) \sim f(x)$

$$p(x) = cf(x)$$

$$Y_i = \frac{f(x_i)}{p(x_i)} = \frac{1}{c}$$

$$\text{Var}(Y) = 0$$

Less variance with better importance sampling


Stratified Sampling

- Estimate subdomains separately

Arvo

Stratified Sampling


- This is still unbiased



$$F_N = \frac{1}{N} \sum_{i=1}^N f(x_i)$$
$$= \frac{1}{N} \sum_{k=1}^M N_k F_k$$

Stratified Sampling

- Less overall variance if less variance in subdomains



$$\text{Var}[F_N] = \frac{1}{N^2} \sum_{k=1}^M N_k \text{Var}[F_k]$$

More Information

- Veach PhD thesis chapter (linked to from website)
- Course Notes (links from website)
 - Mathematical Models for Computer Graphics*, Stanford, Fall 1997
 - State of the Art in Monte Carlo Methods for Realistic Image Synthesis*, Course 29, SIGGRAPH 2001